

# From Coulomb's Law to Gauss's Law

➤ Try to calculate the electric field generated by

\* a point charge **easy**

\* an infinitely long straight wire with evenly distributed charge **hard**

\* a wire loop **only at special locations**

\* a round disk **only at special locations**

\* an infinitely large plane **What??**

\* a solid sphere with evenly distributed charge



➤ Are there other ways to calculate electric field generated from a charge distribution?

➤ Electric field is generated by source charges ➡

are there ways to connect electric field directly with this source charges?

**The answer is YES**

# Electric flux

➤ Electric flux  $\rightarrow$  electric field passing through a given area

➤ For a uniform electric field  $\mathbf{E}$  passing through area  $A$   $\rightarrow$  electric flux is defined as

$$\Phi_E = EA \cos \theta$$




$\theta$   $\rightarrow$  angle between the electric field direction and a line drawn perpendicular to area

➤ Flux can be written equivalently as

$$\Phi_E = E_{\perp} A = EA_{\perp} = EA \cos \theta$$


$E_{\perp} = E \cos \theta$   $\rightarrow$  component of  $\mathbf{E}$  perpendicular to area

$A_{\perp} = A \cos \theta$   $\rightarrow$  projection of area  $A$  perpendicular to field  $\mathbf{E}$




# Electric flux

- \* Electric flux can be interpreted in terms of field lines
- \* Recall that field lines can always be drawn so that number ( $N$ ) passing through unit area perpendicular to field ( $A_{\perp}$ ) is proportional to magnitude of field ( $E$ )



# Gauss Law

- Gauss's law involves the total flux through a closed surface → a surface of any shape that encloses a volume of space



- For any such surface → we divide the surface up into many tiny areas  $\Delta A_1, \Delta A_2, \Delta A_3, \dots$ , and so on
- We make the division so that each  $\Delta A$  is small enough that it can be considered flat and so that the electric field can be considered constant through each  $\Delta A$
- Then the total flux through the entire surface is the sum over all the individual fluxes through each of tiny areas

$$\begin{aligned}\Phi_E &= E_1 \Delta A_1 \cos \theta_1 + E_2 \Delta A_2 \cos \theta_2 + \dots + E_N \Delta A_N \cos \theta_N \\ &= \sum_{j=1}^N E_j \Delta A_j \cos \theta_j = \sum_{j=1}^N E_{\perp j} \Delta A_j = \sum E_{\perp} \Delta A\end{aligned}$$

# Gauss Law

- Number of field lines starting on a positive charge or ending on a negative charge is proportional to magnitude of charge
- Hence  $\blacktriangleright$  the net number of lines  $N$  pointing out of any closed surface (number of lines pointing out minus the number pointing in) must be proportional to the net charge enclosed by the surface  $Q_{\text{encl}}$
- But the net number of lines  $N$  is proportional to the total flux  $\blacktriangleright$

$$\Phi_E = \sum_{\text{closed surface}} E_{\perp} \Delta A \propto Q_{\text{encl}}$$

- To be consistent with Coulomb's law  $\blacktriangleright$  the proportionality constant is  $\epsilon_0^{-1}$


$$\sum_{\text{closed surface}} E_{\perp} \Delta A = \frac{Q_{\text{encl}}}{\epsilon_0}$$

the sum (  $\sum_{\text{closed surface}}$  ) is over any closed surface and  $Q_{\text{encl}}$  is the net charge enclosed within that surface

# Flux Through a Sphere With a Charge at its Center

- A positive point charge  $q$  is located at the center of a sphere of radius  $r$
- According to Coulomb's Law ➡ magnitude of electric field everywhere on surface of sphere is

$$E = k \frac{q}{r^2}$$




# Flux Through a Sphere With a Charge at its Center

- The field lines are directed radially outwards and are perpendicular to surface at every point

$$\Phi_E = \sum_{\text{sphere}} E_{\perp} \Delta A = \sum_{\text{sphere}} E \Delta A = E \sum_{\text{sphere}} \Delta A = E 4\pi r^2$$

- Combine these two equations we have

$$\Phi_E = E \cdot 4\pi r^2 = \frac{1}{4\pi\epsilon_0} \frac{q}{r^2} 4\pi r^2 = \frac{q}{\epsilon_0}$$





# Flux Through a Cube in an Uniform Electric Field


➤ Field lines that pass through surfaces 1 and 2 perpendicularly and are parallel to other four surfaces

➤ For side 1   $\Delta\Phi_E = -E\ell^2$

➤ For side 2   $\Delta\Phi_E = E\ell^2$


➤ For others sides   $\Delta\Phi_E = 0$

➤ Therefore   $\Phi_E = 0$




# Gaussian Surface & Gauss's Law

- You choose a **closed surface** and call it a Gaussian Surface
- This Gaussian Surface can be any shape
- It may or may not enclose charges
- Gauss's Law states



- **Next flux through any closed surface surrounding a charge  $q$  is given by  $q/\epsilon_0$  and is independent of shape of that surface**

$$\sum_{\text{closed surface}} E_{\perp} \Delta A = \frac{Q_{\text{encl}}}{\epsilon_0}$$



# Applying Gauss Law

To use Gauss law  $\rightarrow$  choose a Gaussian surface over which surface  $\Sigma$  can be simplified and electric field determined

**Take advantage of symmetry**


$\rightarrow$  **Gaussian surface is a surface you choose**

**Remember**

$\rightarrow$  **it does not have to coincide with a real surface**

# Field Due to a Spherically Symmetric Even Charge Distribution

- Field must be different inside ( $r < a$ ) and outside ( $r > a$ ) of sphere
- For  $r > a$  select a sphere as Gaussian surface with radius  $r$  and concentric to original sphere
- Because of this symmetry ➡ electric field direction must be radially along  $r$  and at a given  $r$  ➡ field's magnitude is a constant



Can you write down mathematical expression based on above reasoning?

# Field Due to a Spherically Symmetric Even Charge Distribution

- Field must be different inside ( $r < a$ ) and outside ( $r > a$ ) of sphere
- For  $r > a$  select a sphere as Gaussian surface with radius  $r$  and concentric to original sphere
- Because of this symmetry → electric field direction must be radially along  $r$  and at a given  $r$  → field's magnitude is a constant


$E$  is constant at a given  $r$

$$\Phi_E = \sum_{\text{sphere}} E_{\perp} \Delta A = \sum_{\text{sphere}} E \Delta A \stackrel{\downarrow}{=} E \sum_{\text{sphere}} \Delta A = E \cdot 4\pi r^2 \stackrel{\uparrow}{=} \frac{Q}{\epsilon_0}$$

Gauss Law

$$E = \frac{1}{4\pi\epsilon_0} \frac{Q}{r^2} = k \frac{Q}{r^2}$$

$$\vec{E} = \frac{1}{4\pi\epsilon_0} \frac{Q}{r^2} \hat{r} \quad \leftarrow \text{As if the charge is a point charge } Q$$



# Field Inside Sphere

- For  $r < a$  select a sphere as Gaussian surface
- All arguments are same as for  $r > a$
- The only difference is here  $Q_{\text{encl}} < Q$
- Find out that  $Q_{\text{encl}} = Q(r/a)^3$


How?

$$\Phi_E = \sum_{\text{sphere}} E_{\perp} \Delta A = E \cdot 4\pi r^2 = \frac{Q_{\text{encl}}}{\epsilon_0}$$

$$E = \frac{1}{4\pi\epsilon_0} \frac{Q_{\text{encl}}}{r^2} = k \frac{1}{r^2} \frac{r^3}{a^3} Q = k \frac{Q}{a^3} r$$

$$\vec{E} = \frac{1}{4\pi\epsilon_0} \frac{Q}{a^3} \vec{r}$$

Increase linearly with  $r$  not with  $1/r^2$




$$\rho = \frac{Q}{V} \Rightarrow Q_{\text{encl}} = \rho \cdot \frac{4}{3}\pi r^3$$

# Plot Results (Assume Positive $Q$ )

- Inside sphere  $E$  varies linearly with  $r$


$E$  approaches 0 as  $r$  approaches 0



- Field outside sphere is equivalent to that of a point charge located at center of sphere

# Field at a Distance from a Straight Line of Charge


- \* Select a cylinder as Gaussian surface
- \* Cylinder has a radius of  $r$  and a length of  $\ell$
- \*  $\vec{E}$  is constant in magnitude and parallel to surface  
(direction of a surface is its normal!)  
at every point on curved part of surface (body of cylinder)





# Calculate Flux

- \* Because of this line symmetry, end view illustrates more clearly that field is parallel to curved surface, and constant at a given  $r$ , so flux is  $\Phi_E = E \cdot 2\pi r\ell$




- \* Flux through ends of cylinder is 0 since field is perpendicular to these surfaces

# Electric Field from Gauss Law

$$\Phi_E = \sum_{\text{cylinder}} E_{\perp} \Delta A = \frac{q}{\epsilon_0}$$

$$E \cdot 2\pi r \ell = \frac{\lambda \ell}{\epsilon_0}$$

$$E = \frac{\lambda}{2\pi\epsilon_0 r} = 2k \frac{\lambda}{r}$$



One can change thin wire into a rod as we did in sphere case and find electric field inside & outside of rod

# Field Due to an Infinitely Large Plane of Charge


## ➤ Argument about electric field

Because plane is infinitely large, any point can be treated as center point of plane

so at that point  $\vec{E}$  must be parallel to plane direction (again this is its normal)

and must have same magnitude at all points equidistant from plane


## ➤ Choose Gaussian surface to be a small cylinder whose axis is parallel to plane's direction (third time, this is normal of plane)



# Find Out Flux

➤  $\vec{E}$  is parallel to ends  $\rightarrow$  flux through each end of cylinder is  $EA$  and total flux is  $2EA$

➤  $\vec{E}$  is perpendicular to curved surface direction flux through this surface is 0 because  $\cos(90^\circ) = 0$



# Electric Field from Gauss Law

➤ Total charge in surface is  $Q = \sigma A$

➤ Applying Gauss's law

$$\Phi_E = 2EA = \frac{\sigma A}{\epsilon_0} \implies E = \frac{\sigma}{2\epsilon_0}$$


➤ Note, this does not depend on  $r$  → distance from point of interest to charge plane

**WHY?**

➤ Therefore, field is uniform everywhere


# To Summarize

## 3 Types of Gauss Law Problems




$$E = k \frac{Q}{a^3} r \quad \text{👉} \quad r < a$$

$$E = k \frac{Q}{r^2} \quad \text{👉} \quad r \geq a$$



$$E = 2k \frac{\lambda}{r}$$




$$E = \frac{\sigma}{2\epsilon_0}$$

# Conductors in electrostatic equilibrium

- Electrical conductors contain charges (electrons) that are not bound to any atom and therefore are free to move about within the material
- When there is no net motion of charge within a conductor ➡ the conductor is in electrostatic equilibrium
- A conductor in electrostatic equilibrium has the following properties:
  1. Electric field is zero everywhere inside conductor
  2. If an isolated conductor carries a charge ➡ charge resides on its surface
  3. Electric field just outside a charged conductor is perpendicular to surface of conductor and has a magnitude  $\sigma / \epsilon_0$      $\sigma$  ➡ surface charge density at that point!
  4. On an irregularly shape conductor ➤ surface charge density is greatest at locations where radius of curvature of surface is smallest

# Charge distribution in different volumes





## Property 2: For a charged conductor, charge resides on surface, and field inside conductor is still zero


➤ Charges (have to be the same sign, why?) repel and move away from each other until they reach the surface and can no longer move out: charge resides only on the surface because of Coulomb's Law

➤ Choose a Gaussian surface inside but close to the actual surface

➤ Since there is no net charge inside this Gaussian surface, there is no net flux through it.

➤ Because the Gaussian surface can be any where inside the volume


and as close to the actual surface as desired, the electric field inside this volume is zero anywhere



## Property 3: Field's Magnitude and Directions on Surface

### Direction


- Choose a cylinder as the gaussian surface
- The field must be parallel to the surface (again this is its normal)
- \* If there were an angle ( $\theta \neq 0$ ), then there were a component  $E_{\perp}$  from  $\vec{E}$  and tangent to the surface that would move charges along surface  
Then conductor would not be in equilibrium (no charge motions)



## Property 3: Field's Magnitude and Directions on Surface

### Magnitude

- Choose a Gaussian surface as an infinitesimal cylinder with its axis parallel to conductor surface, as shown in figure
- Net flux through Gaussian surface is that only through flat face outside conductor



\* Field here is parallel to surface

\* Field on all other surfaces of Gaussian cylinder is either perpendicular to that surface, or zero

- Applying Gauss's law, we have

$$\Phi_E = EA = \frac{\sigma A}{\epsilon_0} \rightarrow E = \frac{\sigma}{\epsilon_0}$$

# Another Example : Electric Field Generated by a Conducting Sphere and a Conducting Shell


➤ Charge and dimensions as marked

## Analyze

- System has spherical symmetry, Gauss Law problem type I
- Electric field inside conductors is zero
- There are two other ranges,  $a < r < b$  and  $c < r$  that need to be considered

➤ Arguments for electric field

- Similar to sphere example, because spherical symmetry, electrical field in these two ranges  $a < r < b$  and  $c < r$  is only a function of  $r$ , and goes along radius



# Construct Gaussian Surface & Calculate Flux & Use Gauss Law To Get Electric Field

- $E = 0$  when  $r < a$ , and  $b < r < c$
- Construct a Gaussian sphere with its center coincides with center of inner sphere

➤ When  $a < r < b$

- Flux  $\Phi_E = E \cdot 4\pi r^2$

- Apply Gauss Law  $\Phi_E = \frac{Q}{\epsilon_0}$


$$E = \frac{1}{4\pi\epsilon_0} \frac{Q}{r^2} \quad \text{or} \quad \vec{E} = \frac{1}{4\pi\epsilon_0} \frac{Q}{r^2} \hat{r}$$

➤ When  $c < r$


- Flux  $\Phi_E = E \cdot 4\pi r^2$

- Apply Gauss Law  $\Phi_E = \frac{-2Q + Q}{\epsilon_0}$

$$E = \frac{1}{4\pi\epsilon_0} \frac{-Q}{r^2} \quad \text{or} \quad \vec{E} = \frac{1}{4\pi\epsilon_0} \frac{-Q}{r^2} \hat{r}$$



# Definition of a Faraday Cage



# A Practical conclusion from Gauss's Law


## Faraday's Cage

➤ The field induces charges on the left and right sides of the conductive box



➤ The total electric field inside the box is zero

➤ The presence of the box distorts the Field in adjacent regions



**During a thunderstorm stay in your car!!**

# When a Car is Struck by Lightning

**1** Lightning strikes car frame

**2** Current travels around the outside of the vehicle

**3** Current exits to the ground through the car







---